

HEAVY QUARK PRODUCTION AT HERA IN K_T FACTORIZATION SUPPLEMENTED WITH CCFM EVOLUTION

H. JUNG

*Physics Department, Lund University, Box 118,
S-221 00 Lund, Sweden
E-mail: Hannes.Jung@desy.de*

The application of k_t - factorization, supplemented with the CCFM small- x evolution equation, to heavy quark production is discussed. Differential cross sections of $b\bar{b}$ production and also inelastic J/ψ production as measured at HERA are compared to the hadron level CCFM Monte Carlo generator CASCADE, using the unintegrated gluon density obtained within the CCFM evolution approach from a fit to HERA F_2 data.

1 Introduction

The calculation of inclusive quantities, like the structure function $F_2(x, Q^2)$ at HERA, performed in NLO QCD is in perfect agreement with the measurements. But in exclusive quantities like jet or heavy quark production large so-called K -factors (normalization factors)^{1,2} are needed to bring the NLO calculations close to the data.

At large energies (small x), the evolution of parton densities proceeds over a large region in rapidity $\Delta y \sim \log(1/x)$ and effects of finite transverse momenta of the partons may become increasingly important. Cross sections can then be k_t - factorized³ into an off-shell (k_t dependent) partonic cross section $\hat{\sigma}(\frac{x}{z}, k_t)$ and a k_t - unintegrated parton density function $\mathcal{F}(z, k_t)$:

$$\sigma = \int \frac{dz}{z} d^2 k_t \hat{\sigma}\left(\frac{x}{z}, k_t\right) \mathcal{F}(z, k_t) \quad (1)$$

The unintegrated gluon density $\mathcal{F}(z, k_t)$ is described by the BFKL evolution equation in the region of asymptotically large energies (small x). An appropriate description valid for both small and large x is given by the CCFM evolution equation^{4,5,6,7}, resulting in an unintegrated gluon density $\mathcal{A}(x, k_t, \bar{q})$, which is a function also of the additional evolution scale $\bar{q}$.

Catani^{8,9} argues that by explicitly carrying out the k_t integration in eq.(1) one can obtain a form fully consistent with collinear factorization: the coefficient functions and also the DGLAP splitting functions are no longer evaluated in fixed order perturbation theory but supplemented with the all-order resum-

mation of the $\alpha_s \log 1/x$ contribution at small x . This all-loop resummation shows up in a so-called *non-Sudakov* form factor Δ_{ns} for CCFM.

In this paper the k_t factorization approach is discussed with respect to resolved photon structure and next-to-leading corrections in the collinear approach. Then calculations for $b\bar{b}$ and inelastic J/ψ production at HERA are presented, using the unintegrated gluon density obtained previously¹⁰ from a CCFM fit to the HERA structure function $F_2(x, Q^2)$

2 k_t -factorization versus higher order processes in collinear factorization




Figure 1. Diagrammatic representation of LO, NLO and resolved photon processes in the collinear approach compared to the k_t -factorization approach.

The diagrams for LO, NLO and resolved photon processes in the collinear approach are shown schematically in Fig. 1. In collinear factorization, the incoming parton is on-mass-shell and has vanishing transverse momentum k_t , whereas in k_t -factorization the partons entering the hard scattering matrix element are free to be off-mass-shell. The advantage of the k_t -factorization approach becomes visible, when additional hard gluon radiation to a $2 \rightarrow 2$ process like $\gamma g \rightarrow Q\bar{Q}$ is considered. If the transverse momentum p_{tg} of the additional gluon is of the order of that of the quarks, then in the collinear approach the full $\mathcal{O}(\alpha_s^2)$ matrix element for $2 \rightarrow 3$ has to be calculated. In k_t -factorization such processes are naturally included, even if only the LO α_s off-shell matrix element is used, since the k_t of the incoming gluon is not restricted from above, and therefore can acquire a virtuality similar to the ones in a complete fixed order calculation. In Fig. 1 the basic ideas are shown schematically. Not only does k_t -factorization include (at least parts of) NLO diagrams, it also includes diagrams of the resolved photon type¹¹, with the natural transition from real to virtual photons.

The $\mathcal{O}(\alpha_s)$ matrix element in k_t -factorization includes the $\mathcal{O}(\alpha_s)$ matrix element of collinear factorization but in addition also higher order contributions because the incoming gluon is off-shell and the unintegrated gluon density resums parts of the virtual corrections (Fig. 2).




Figure 2. Schematic diagrams for k_t -factorization: left is shown the one-loop correction to the Born diagram for photo-production right is shown the all-loop improved correction with the factorized structure function $\mathcal{F}(x, k_t^2)$

The unintegrated gluon density $x\mathcal{A}(x, k_t^2, \bar{q})$ ^{10,13,14}, was obtained in the framework of k_t factorization using the CCFM evolution equation to fit¹⁰ the structure function $F_2(x, Q^2)$. This gluon density $x\mathcal{A}(x, k_t^2, \bar{q})$ is used in the hadron level Monte Carlo generator CASCADE¹⁵ for any comparison with measurements.

3 Heavy quark production at HERA

The prediction of CASCADE for the total $b\bar{b}$ cross section¹⁰ was compared to the extrapolated measurements of the H1¹⁶ and ZEUS¹⁷ experiments at HERA. Since CASCADE generates full hadron-level events, a direct comparison with measurements can be done, before extrapolating the measurement over the full phase space to the total $b\bar{b}$ cross section. It has been shown¹³, that the visible $b\bar{b}$ cross section as measured by the ZEUS experiment¹⁷ is in perfect agreement with the predictions from CASCADE. However, large uncertainties come from the extrapolation of the measured cross section to the total $b\bar{b}$ cross section. If the extrapolation is performed with CASCADE (or any other program of the Lund family), approximate agreement with the NLO prediction is also achieved in the case of the ZEUS measurement¹⁷.

The prediction of CASCADE has also been compared¹³ to the measurement of H1¹⁶, where it was also shown that the experimental results from H1 and




Figure 3. The fractional energy x_g of the gluon of the process $\gamma g^* \rightarrow b\bar{b}$ satisfying the kinematic selection of ZEUS¹².

ZEUS are different by a factor ~ 2 . It has been checked, that the difference between the experiments is not due to different kinematics.




Figure 4. Differential bottom cross section ¹² as a function of the muon pseudo-rapidity (transverse momentum) for event with two jets and a muon, compared to CASCADE and PYTHIA (with and w/o heavy quark excitation).

In Fig. 4 the differential cross section of μ 's coming from b -quark decays as measured by ZEUS¹² is compared with the prediction of CASCADE. Again good agreement is observed, except in the region of large η^μ . However this region is dominated by large $x_g \gtrsim 0.03$ (see Fig. 3), where the application of k_t -factorization is questionable and also where the unintegrated gluon density is not at all constrained by the fit to F_2 .




Figure 5. The cross section of $\gamma p \rightarrow J/\psi + X$ as a function of the transverse momentum (a) and of the fractional momentum z (b) of the J/ψ as measured by H1¹⁸ compared to the predictions of CASCADE and EPJPSI.

The cross section for $\gamma p \rightarrow J/\psi + X$ as a function of the transverse momen-

tum of the J/ψ shows a significant deviation from the leading order color singlet model prediction (as implemented in EPJPSI¹⁹) in collinear factorization. In Fig. 5, the preliminary H1 measurement¹⁸ is compared to the prediction of EPJPSI¹⁹. In collinear factorization, the harder transverse momentum spectrum is interpreted as a signal for significant next-to-leading order corrections. Also shown in Fig. 5a is the prediction of CASCADE using the same CCFM unintegrated gluon density as before together with the k_t -factorized matrix element²⁰ for $\gamma g^* \rightarrow J/\psi g$. The inelastic J/ψ photoproduction cross section ($z < 0.9$) as a function of p_T is nicely described by CASCADE. Especially the large transverse momentum part is explained as additional hard initial state QCD radiation. The same is also observed in full NLO calculations, and it shows again the advantage of the k_t -factorization in simulating a large part of the NLO correction of the collinear approach, due to the non zero virtuality of the incoming gluon. The distribution in the fractional momentum z of the J/ψ is also reasonably well described (Fig. 5b).

4 Conclusion

The application of the k_t -factorization approach to heavy quark production at HERA has been discussed. It is shown, that the visible $b\bar{b}$ production cross section as measured at ZEUS is nicely reproduced by CASCADE, using the CCFM evolved unintegrated gluon density obtained from a fit to $F_2(x, Q^2)$. It was pointed out, that the extrapolation from the measured to the total $b\bar{b}$ cross section contains large model dependencies.

The differential cross sections for inelastic J/ψ photoproduction can be reasonably well described with CASCADE, both in terms of shape and normalization. The large transverse momentum tail is a direct signal for additional hard initial state QCD radiation.

It is the advantage of the k_t -factorization approach that important parts of NLO and even NNLO contributions are consistently included due to the off-shell gluons, which enter into the hard scattering process.

Acknowledgments

I want to thank the organizers M. Kienzle and M. Wadhwa for this very nice workshop. I am also grateful to S. Frixione for the invitation to this workshop and his interest in k_t -factorization. Many thanks also go to S. Baranov and N. Zotov for interesting discussions and our fruitful collaboration. I am also grateful to L. Gladilin, B. Naroska and J. Whitmore for careful reading of the manuscript. All thanks for the great times with Antje.

References

1. CDF Collaboration; F. Abe et al., *Phys. Rev. D* **55** (1997) 2546.
2. D0 Collaboration; B. Abbott et al., *Phys. Lett. B* **487** (2000) 264.
3. S. Catani, M. Ciafaloni, F. Hautmann, *Nucl. Phys. B* **366** (1991) 135.
4. M. Ciafaloni, *Nucl. Phys. B* **296** (1988) 49.
5. S. Catani, F. Fiorani, G. Marchesini, *Phys. Lett. B* **234** (1990) 339.
6. S. Catani, F. Fiorani, G. Marchesini, *Nucl. Phys. B* **336** (1990) 18.
7. G. Marchesini, *Nucl. Phys. B* **445** (1995) 49.
8. S. Catani, k_t -factorisation and perturbative invariants at small x , in *Proceedings of the International Workshop on Deep Inelastic Scattering, DIS 96* (Rome, Italy, 15-19 April, 1996), hep-ph/9608310.
9. S. Catani, Aspects of QCD, from the Tevatron to LHC, in *Proceedings of the International Workshop Physics at TeV Colliders* (Les Houches, France, 8-18 June, 1999), hep-ph/0005233.
10. H. Jung, G. Salam, *Eur. Phys. J. C* **19** (2001) 351, hep-ph/0012143.
11. S. Baranov, N. Zotov, *Phys. Lett. B* **491** (2000) 111.
12. ZEUS Collaboration; M. Derrick et al., Beauty photoproduction in the muon semi-leptonic decay mode at HERA, in *Contributed paper 496 to IECHEP 2001, Budapest, Hungary* (2001).
13. H. Jung, Unintegrated parton densities applied to heavy quark production in the CCFM approach, in *Proceedings of the Rinberg workshop on "New trends in HERA physics", Ringberg Castle, Tegernsee, Germany.* (2001), hep-ph/0109146.
14. H. Jung, *submitted to Phys. Rev. D* (2001), DESY-01-136, hep-ph/0110034.
15. H. Jung, *The CCFM Monte Carlo generator CASCADE for lepton - proton and proton - proton collisions*, Lund University, accepted by *Comp. Phys. Comm.*, 2001, hep-ph/0109102, DESY 01-114, <http://www.quark.lu.se/~hannes/cascade/>.
16. H1 Collaboration; C. Adloff et al., *Phys. Lett. B* **467** (1999) 156, and erratum *ibid*.
17. ZEUS Collaboration; J. Breitweg et al., *Eur. Phys. J. C* **18** (2001) 625.
18. H1 Collaboration; C. Adloff et al., Inelastic photoproduction of J/ψ and $\psi(2s)$ at H1, in *Contributed paper 157aj to IECHEP 1999, Tampere, Finland* (1999).
19. H. Jung, *EPJPSI: A Monte Carlo generator for J/ψ mesons in high energy $\gamma - p$, $e - p$ and $p - \bar{p}$ collisions, version 3.3*, Lund University, 1995, <http://www-h1.desy.de/~jung/epjpsi.html>.
20. V. Saleev, N. Zotov, *Mod. Phys. Lett. A* **9** (1994) 151.